

THE SANDWICH THEOREM FOR SUBLINEAR AND SUPERLINEAR FUNCTIONALS

A.T. DIAB, S. I. NADA, D. L. FEARNLEY

ABSTRACT. The Hahn-Banach theorem is an extension theorem for linear functionals which preserves certain properties. Specifically, if a linear functional is defined on a subspace of a real vector space which is dominated by a sublinear functional on the entire space, then this functional can be extended to a linear functional on the entire space which is still dominated by a sublinear functional. In this paper, we generalize this result to show that a linear functional defined on a subspace of a real vector space which is dominated by a sublinear functional and also dominates a superlinear functional on the entire space can be extended to a linear functional on the entire space which is also dominated by a sublinear functional and dominates a superlinear functional.

1. INTRODUCTION

Extensions of linear functionals have been studied in locally convex spaces by several authors. In fact, the importance of the Hahn-Banach theorem arises from its wide variety of applications, including complex and functional analysis and thermodynamics. The Hahn-Banach theorem has important implications for convex sets, and is the foundation for an effective treatment of optimization. It is also used to solve many problems in linear algebra, conic duality theory, the minimax theorem, piecewise approximation of convex functionals, extensions of positive linear functionals, and other results from modern control [1,5,8]. The Hahn-Banach theorem was proved by Hans Hahn (1879- 1934) in 1927 and later by Stefan Banach (1892-1945) in 1929. It states that if E is a real vector space, M is a subspace of E and S is a sublinear functional on E , and f_0 is a linear functional on M such that $f_0(x) \leq S(x)$ for every $x \in M$ (or equivalently, $-S(-x) \leq f_0(x) \leq S(x)$), then there exists a linear functional L on E such that $L|_M = f_0$ and $L(x) \leq S(x)$ for every $x \in E$ (or equivalently, $-S(-x) \leq L(x) \leq S(x)$). We will show that if P is a superlinear functional on E satisfying the condition

2000 *Mathematics Subject Classification.* 46A22 Extension linear functionals, The Hahn - Banach theorem, The classical Hahn - Banach theorem, Sublinear functionals and superlinear functionals.

that $P(x) \leq f_0(x) \leq S(x)$ for every $x \in M$ such that $f_0(x) \leq T(x)$ for every $x \in M$, where $T(x) = \inf_{y \in E} \{S(x+y) - P(y)\}$ for every $x \in E$, then there exists a linear functional L on E such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in E$. We will refer to this as the sandwich theorem for a sublinear and superlinear functionals. Note that if we take $P(x) = -S(-x)$ for every $x \in E$, then we obtain the Hahn-Banach theorem. As an application of the Hahn-Banach theorem we have the following theorem: There exists a linear functional L on E for every sublinear functional S on E such that $L(x) \leq S(x)$ for every $x \in E$ (we refer to this as the classical Hahn-Banach theorem). We will also use the sandwich theorem for a sublinear and superlinear functionals to show that, for every a sublinear functional S on E and for every superlinear functional P on E , if $P(x) \leq S(x)$ for every $x \in E$ then there exists a linear functional L on E such that $P(x) \leq L(x) \leq S(x)$ for every $x \in E$. Note that if we take $P(x) = -S(-x)$ for every $x \in E$, then this result implies the classical Hahn-Banach theorem.

2. RELATIONS BETWEEN SUBLINEAR AND SUPERLINEAR FUNCTIONALS DEFINED ON A REAL LINEAR SPACE

In this section we present some basic definitions and results from functional analysis concerning sublinear and superlinear functionals. In the following definitions we suppose that E is a real vector space [6,7,9].

Definition 2.1. Let E be a real vector space, M be a subspace of E and f_0 be a linear functional on M . Then a linear functional L on E is called an *extension functional* of f_0 if $L(x) = f_0(x)$ for every $x \in M$ (denoted $L|_M = f_0$).

Definition 2.2. The functional $S : E \rightarrow \mathbb{R}$ is called *sublinear* if it possesses the following properties:

- (1) $S(x+y) \leq S(x) + S(y)$ for every $x, y \in E$ (i.e., S is subadditive),
- (2) $S(\alpha x) = \alpha S(x)$ for every $x \in E$ and $\alpha > 0$ (i.e., S is positively homogeneous). The set of all sublinear functionals on E is denoted by $Subl(E)$.

Definition 2.3. The functional $P : E \rightarrow \mathbb{R}$ is called *superlinear* if it possesses the following properties:

(1) $P(x+y) \geq P(x) + P(y)$ for every $x, y \in E$ (i.e., P is superadditive),
 (2) $P(\alpha x) = \alpha P(x)$ for every $x \in E$ and $\alpha > 0$ (i.e., P is positively homogeneous). The set of all superlinear functionals on E is denoted by $Supl(E)$.

Note that $Lin(E) = Subl(E) \cap Supl(E)$, where $Lin(E)$ is the set of all linear functionals on E . Also, it is easy to verify that if $E \equiv \mathbb{R}^n$, $S(x) = \sqrt{\sum_{i=1}^n x_i^2} = \|x\|$ and $P(x) = -\sqrt{\sum_{i=1}^{n-1} x_i^2} + x_n$ for every $x \in \mathbb{R}^n$, then $S \in Subl(E)$ and $P \in Supl(E)$. Also, $S \in Subl(E)$ if and only if $-S \in Supl(E)$.

Throughout the this paper we will use the symbols S and P to denote sublinear and superlinear functionals on the real vector space E .

Lemma 2.1. (i) $S(\underline{0}) = 0$ and $P(\underline{0}) = 0$, where $\underline{0}$ is the zero vector of the real vector space E .

(ii) $-S(-x) \leq S(x)$ and $-P(-x) \geq P(x)$ for every $x \in E$.

(iii) A functional f on E is linear if and only if it is additive and positively homogeneous.

Proof. (i) Since sublinear functionals are positively homogeneous, $S(\underline{0}) = S(2\underline{0}) = 2.S(\underline{0} + \underline{0}) = S(\underline{0}) + S(\underline{0})$. Therefore $S(\underline{0}) = 0$. Similarly, $P(\underline{0}) = 0$.

(ii) Since $0 = S(\underline{0}) = S(-x + x) \leq S(-x) + S(x)$ for every $x \in E$, then $-S(-x) \leq S(x)$. Also, $0 = P(\underline{0}) = P(-x + x) \geq P(-x) + P(x)$ for every $x \in E$, so $-P(-x) \geq P(x)$ for every $x \in E$.

(iii) By definition, linear functionals are additive and positively homogeneous. Let f be an additive positively homogeneous functional on E . It remains to show that $f(\alpha x) = \alpha f(x)$ for every $x \in E$ and $\alpha < 0$. Since f is additive, $f(\underline{0} + \underline{0}) = f(\underline{0}) + f(\underline{0})$. Thus, $f(\underline{0}) = 0$, and consequently $f(-x) = -f(x)$ for every $x \in E$. Let $x \in E$ and $\alpha = -t$, where $t > 0$. Then $f(\alpha x) = f(-tx) = tf(-x) = -tf(x) = \alpha f(x)$. Thus f is homogeneous and f is linear.

Lemma 2.2. If $S(x) \leq P(x)$ for every $x \in E$, then P and S are linear functionals on E . Moreover $S \equiv P$ on E .

Proof. Since S is sublinear and P is superlinear, then from (ii) in Lemma 2.1, we obtain that $-S(-x) \leq S(x)$ and $-P(-x) \geq P(x)$ for every $x \in E$. So, $S(-x) \leq P(-x) \leq -P(x) \leq -S(x)$. Thus,

$S(-x) = -S(x)$ for every $x \in E$. For all $x, y \in E$, $S(x + y) = -S(-x - y) \geq -S(-x) - S(-y)$. Hence, S is both subadditive and superadditive and is therefore additive. Since S is positively homogeneous and additive, from (iii) in Lemma 2.1 we can conclude that S is linear. By a similar argument, P is linear. Finally, from (ii) in Lemma 2.1 we see that $-S(x) \leq S(-x) \leq P(-x) \leq -P(x)$. Hence, $P(x) \leq S(x)$ for every $x \in E$. Therefore $P(x) = S(x)$ for every $x \in E$.

Lemma 2.3. *If f is a homogeneous functional on E and g is an arbitrary functional on E , then the following statements are equivalent:*

- (i) $f(x) \leq g(x)$ for every $x \in E$,
- (ii) $-g(-x) \leq f(x) \leq g(x)$ for every $x \in E$.

Proof. Clearly, (ii) implies (i). Let $f(x) \leq g(x)$ for every $x \in E$. Then $f(-x) \leq g(-x)$ and consequently $-f(x) = f(-x) \leq g(-x)$ for every $x \in E$. Therefore $-g(-x) \leq f(x)$ and hence $-g(-x) \leq f(x) \leq g(x)$.

Corollary 2.1. *If f is a linear functional on E and $f(x) \leq S(x)$ for every $x \in E$, where S is a sublinear functional on E , then $-S(-x) \leq f(x) \leq S(x)$ for every $x \in E$.*

Proof. The proof follows directly from Lemma 2.3 and the fact that every linear functional is homogeneous.

Lemma 2.4. *Let $f \in \text{Lin}(M)$, and let $g, h : M \rightarrow \mathbb{R}$, where M is a subspace of E , $g(\underline{0}) = h(\underline{0}) = 0$, and $g(x) \leq h(x)$ for every $x \in M$. Then the following are equivalent:*

- (i) $g(x) \leq f(x) \leq h(x)$ for every $x \in M$,
- (ii) $f(x) \leq h(x + y) - g(y)$ for all $x, y \in M$.

Proof. First, we assume that $g(x) \leq f(x) \leq h(x)$ for every $x \in M$, and we want to show that $f(x) \leq h(x + y) - g(y)$ for every $x, y \in M$. Since $g(x + y) \leq f(x + y) \leq h(x + y)$, and $-h(y) \leq -f(y) \leq -g(y)$, it follows that $g(x + y) - h(y) \leq f(x + y) - f(y) \leq h(x + y) - g(y)$. Since $f \in \text{Lin}(M)$, then $g(x + y) - h(y) \leq f(x) \leq h(x + y) - g(y)$. Next, assume that $f(x) \leq h(x + y) - g(y)$ for every $x, y \in M$. Setting $y = \underline{0}$ we see that $f(x) \leq h(x)$. By setting $x = -y$, we obtain $f(-y) \leq -g(y)$, so $g(y) \leq f(y)$. Therefore, $g(x) \leq f(x) \leq h(x)$ for every $x \in M$.

3. THE HAHN-BANACH THEOREM AND THE CLASSICAL HAHN-BANACH THEOREM

Theorem 3.1. *(The Hahn-Banach theorem for sublinear functionals)*

Let E be a real vector space, M be a subspace of E and $S \in \text{Subl}(E)$. Furthermore, let $f_0 \in \text{Lin}(M)$ such that $f_0(x) \leq S(x)$ for every $x \in M$ (or equivalently, $-S(-x) \leq f_0(x) \leq S(x)$), then there exists a functional $L \in \text{Lin}(E)$ such that $L|_M = f_0$ and $L(x) \leq S(x)$ for every $x \in E$ (or equivalently, $-S(-x) \leq f_0(x) \leq S(x)$) [7, 3].

The following theorem is the famous application of the Hahn-Banach theorem and some mathematicians, like Kelly-Namioka[2], Rudin [7], Koing [4], Simons [8,9] and others have proved it. In the following, we introduce a simple and short proof for it.

Theorem 3.2. *(The classical Hahn-Banach theorem for sublinear functionals)* Let E be a real vector space and $S \in \text{Subl}(E)$, then there exists a functional $L \in \text{Lin}(E)$ such that $L(x) \leq S(x)$ for every $x \in E$.

Proof. If $S(x) = 0$ for every $x \in E$, then the result follows immediately by taking $L \equiv 0$ on E . Assume that $S \not\equiv 0$ on E . Let $x_0 \in E$, let $M = \{\alpha x_0 : \alpha \in \mathbb{R}\}$, and define $f_0(\alpha x_0) = \alpha S(x_0)$ for every $x \in M$. We have two cases:

Case 1: $M = \{0\}$. Since $0 = f_0(0) = S(0)$, $f_0(x) \leq S(x)$ for every $x \in M$. By the Hahn-Banach theorem (or Theorem 3.1), there exists a linear functional L on E such that $L|_M = f_0$ and $L(x) \leq S(x)$ for every $x \in E$.

Case 2: $M \neq \{0\}$. Let $x = \alpha x_0 \in M$. If $\alpha \geq 0$ then $f_0(x) = f_0(\alpha x_0) = \alpha f_0(x_0) \leq \alpha S(x_0) = S(x)$. If $\alpha < 0$ then let $\alpha = -t$, where $t > 0$. Then $f_0(x) = f_0(\alpha x_0) = \alpha S(x_0) = -tS(x_0) = -S(tx_0) = -S(-\alpha x_0) = -S(-x) \leq S(x)$ for every $x \in M$. Hence, $f_0(x) \leq S(x)$ for every $x \in M$. Thus, from the extension form of the Hahn-Banach theorem, there exists a linear functional L on E such that $L|_M = f_0$ and $L(x) \leq S(x)$ for every $x \in E$.

In the following Theorems, we mention a similar fact for superlinear functionals and we call it the extension form of the Hahn-Banach theorem for superlinear functionals and the classical Hahn-Banach theorem for superlinear functionals [6]. The proofs of Theorems 3.3 and 3.4 can be found in [6].

Theorem 3.3. *(The extension form of Hahn-Banach theorem for superlinear functionals)*

Suppose that $f_0 \in \text{Lin}(M)$, where M is a subspace of E such that $P(x) \leq f_0(x)$ for every $x \in M$, where $P \in \text{Supl}(E)$. Then there exists a functional $L \in \text{Lin}(E)$ such that $L|_M = f_0$ and $P(x) \leq L(x)$ for every $x \in E$.

Theorem 3.4. (The classical Hahn-Banach theorem for superlinear functionals)

Let E be a real vector space and $P \in \text{Supl}(E)$, then there exists a functional $L \in \text{Lin}(E)$ such that $P(x) \leq L(x)$ for every $x \in E$.

4. THE SANDWICH THEOREM FOR SUBLINEAR AND SUPERLINEAR FUNCTIONALS

In this section we demonstrate three different approaches to proving that, if $S \in \text{Subl}(E)$, $P \in \text{Supl}(E)$ and $f_0 \in \text{Lin}(M)$, where M is a subspace of E , and $P(x) \leq f_0(x) \leq S(x)$ for each $x \in M$, then there is an extension functional $L \in \text{Lin}(E)$ so that $P(x) \leq L(x) \leq S(x)$ for each $x \in E$. Throughout this section, we let $T(x) = \inf_{y \in E} \{S(x + y) - P(y)\}$ for every $x \in E$.

Lemma 4.1. Let f_0 be a linear functional of a subspace M of E such that: $f_0(x) \leq T(x)$ for every $x \in M$ (4.1). Then the following conditions are satisfied:

$P(x) \leq S(x)$ for every $x \in E$ (4.2),

$P(x) \leq f_0(x) \leq S(x)$ for every $x \in M$ (4.3).

Proof. From the inequality (4.1), we obtain that $f_0(x) \leq S(x + y) - P(y)$ for every $y \in E$ and $x \in M$. Consequently, by setting $x = \underline{0}$, we see that $P(y) \leq S(y)$ for every $y \in M$. By setting $y = \underline{0}$, we see that $f_0(x) \leq S(x)$ for every $x \in M$, and by setting $y = -x$, we obtain that $f_0(-y) \leq -P(y)$ or $P(x) \leq f_0(x)$ for every $x \in M$. Hence $P(x) \leq f_0(x) \leq S(x)$ for every $x \in M$.

The following example shows that the converse of the above lemma is false.

Example 4.1. Let $E = \mathbb{R}^2$, $P(x) = -|x_1| + x_2$, $S(x) = \sqrt{x_1^2 + x_2^2}$, where $x = (x_1, x_2)$ and $M = \{(x_1, 0) : x_1 \in \mathbb{R}\}$ is an one dimensional subspace of E . Suppose that $f_0 : M \rightarrow \mathbb{R}$ is defined as $f_0(x) = x_1$ for

every $x = (x_1, 0) \in M$. Then inequalities (4.2) and (4.3) are satisfied but (4.1) is not.

Solution. It is easy to verify that $P \in \text{Supl}(E)$, $S \in \text{Subl}(E)$ and f_0 is a linear functional on M . Let $x = (x_1, x_2) \in E$. Then $S(x) = \sqrt{x_1^2 + x_2^2} \geq x_2 \geq -|x_1| + x_2 = P(x)$. Also, if $x = (x_1, 0) \in M$ then $P(x) = -|x_1| \leq x_1 = f_0(x) \leq |x_1| \leq \sqrt{x_1^2 + x_2^2} = S(x)$, so $P(x) \leq f_0(x) \leq S(x)$ for every $x \in M$. Finally, suppose that inequality (4.1) is true. Then, for each $x \in M$, $f_0(x) \leq S(x+y) - P(y)$ for every $y \in E$. Let $x = (x_1, 0) \in M$ where $x_1 > 0$. Then $x_1 \leq \sqrt{(x_1 + y_1)^2 + y_2^2} + |y_1| - y_2$ for every $y = (y_1, y_2) \in E$. In particular, if $y = (0, y_2)$ and $y_2 > 0$ then this inequality becomes $x_1 \leq \sqrt{x_1^2 + y_2^2} - y_2 = \frac{(\sqrt{x_1^2 + y_2^2} - y_2)(\sqrt{x_1^2 + y_2^2} + y_2)}{(\sqrt{x_1^2 + y_2^2} + y_2)} = \frac{x_1^2}{\sqrt{x_1^2 + y_2^2} + y_2}$, which approaches 0 as $y_2 \rightarrow \infty$. But this implies that $x_1 \leq 0$, a contradiction.

Lemma 4.2. *If $P(x) = -S(-x)$ for every $x \in E$, then $T(x) = S(x)$ for every $x \in E$.*

Proof. From the definition of T , we conclude that $T(x) \leq S(x+y) - P(y)$ for all $x, y \in E$. Thus $T(x) \leq S(x)$. Also, $S(x) \leq S(x+y) + S(-y) = S(x+y) - P(y)$ for every $y \in E$, so $S(x) \leq T(x)$. Therefore $T(x) = S(x)$ for every $x \in E$, as required.

Corollary 4.1. *If $P(x) = -S(-x)$ for every $x \in E$, then conditions (4.2) and (4.3) are satisfied if and only if condition (4.1) is satisfied.*

Proof. Sufficiency follows from Lemma 4.1. Conversely, if conditions (4.2) and (4.3) are satisfied then condition (4.1) follows from Lemma 4.2.

Lemma 4.3. *Let $P(x) \leq S(x)$ for every $x \in E$. Then $P(x) \leq T(x) \leq S(x)$ for every $x \in E$, and T is a sublinear functional. Furthermore, if L is a linear functional on E such that $P(x) \leq L(x) \leq S(x)$ for every $x \in E$ then $L(x) \leq T(x)$ for every $x \in E$.*

Proof. Since $P(u) \leq L(u)$, it follows that $L(u) \leq -P(-u)$. Hence, from the linearity of L we see that $L(u+v) \leq S(v) - P(-u)$ for every $u, v \in E$. Setting $u+v = x$ and $u = -y$, we obtain $L(x) \leq S(x+y) - P(y)$. Taking the infimum over all $y \in E$, we see that $L(x) \leq T(x)$ for every $x \in E$. We next prove that $T : E \rightarrow \mathbb{R}$ is

sublinear. Since $P(y) \leq S(y) \leq S(x+y) + S(-x)$, it follows that $-S(-x) \leq S(x+y) - P(y)$. Taking the infimum over all $y \in E$, we see that $T(x) \geq -S(-x)$, so $T : E \rightarrow \mathbb{R}$. It follows that T is positively homogeneous since S and P are positively homogeneous. Thus, to prove that T is sublinear it remains to show that T is subadditive. Let $u, v \in E$. Then $T(u+v) = \inf_{y+z \in E} \{S(u+v+y+z) - P(y+z)\} \leq \inf_{y \in E} \{S(u+y) - P(y)\} + \inf_{z \in E} \{S(v+z) - P(z)\} = T(u) + T(v)$. Thus T is subadditive and, consequently, sublinear. Since $T(x) \leq S(x+y) - P(y)$, it follows that $T(x) \leq S(x+\underline{0}) - P(\underline{0})$, so $T(x) \leq S(x)$ for every $x \in E$. Also, $T(-y) \leq S(\underline{0}) - P(y)$, so $T(-y) \leq -P(y)$ and $-T(y) \leq T(-y) \leq -P(y)$, and hence $P(y) \leq T(y)$ for every $y \in E$. Therefore $P(x) \leq T(x) \leq S(x)$ for every $x \in E$, and the proof is complete.

In the following theorem we introduce the first method for proving the statement above, which we call the sandwich theorem for sublinear and superlinear functionals. Throughout this section, we assume that f_0 is a linear functional on M , where M is a subspace of E . As before, $S \in \text{Subl}(E)$ and $P \in \text{Supl}(E)$.

Theorem 4.1. *There exists an extension linear functional L on E of the linear functional f_0 on M such that $P(x) \leq L(x) \leq S(x)$ for every $x \in E$ if and only if $f_0(x) \leq T(x)$ for every $x \in M$.*

Proof. From Lemma 4.3 we know that $T \in \text{Subl}(E)$ and $P(x) \leq T(x) \leq S(x)$ for every $x \in E$. Also by lemma 4.3, if $P(x) \leq f_0(x) \leq S(x)$ for every $x \in M$ then $f_0(x) \leq T(x)$ for every $x \in M$ since f_0 is a linear functional on M . Next, assume that $f_0(x) \leq T(x)$ for every $x \in M$. By lemma 4.1, $P(x) \leq S(x)$ for all $x \in E$ and $P(x) \leq f_0(x) \leq S(x)$ for all $x \in M$. By Theorem 3.3 there is an extension linear functional L on E such that $f_0(x) = L(x)$ for every $x \in M$ and $P(x) \leq L(x)$ for every $x \in E$. By lemma 4.3 $L(x) \leq T(x) \leq S(x)$ for every $x \in E$. Hence, it follows that $P(x) \leq L(x) \leq S(x)$ for every $x \in E$.

The following corollary shows that if we let $P(x) = -S(-x)$ for every $x \in E$, then we obtain the Hahn-Banach theorem.

Corollary 4.2. *If $P(x) = -S(-x)$ for every $x \in E$ and f_0 is a linear functional on M such that $f_0(x) \leq S(x)$ for every $x \in M$. Then there exists a linear functional L on E such that $L|_M = f_0$ and $L(x) \leq S(x)$ for every $x \in E$.*

Proof. The proof follows directly from theorem 4.1 and corollary 4.1, as required.

We now introduce the second proof of the Sandwich Theorem for Superlinear and Sublinear Functionals.

Lemma 4.4. *Let $x_0 \notin M$, $a = \sup_{y \in M} \{f_0(y) - S(y - x_0)\}$, $b = \inf_{y \in M} \{-f_0(y) + S(y + x_0)\}$, $c = \inf_{y \in M} \{f_0(y) - P(y - x_0)\}$ and $d = \sup_{y \in M} \{-f_0(y) + P(y + x_0)\}$, and assume that $a \leq b$ and $d \leq c$. Then the following statements are equivalent:*

- (i) $f_0(x) \leq S(x + y) - P(y)$ for every $x \in M$ and $y \in E$.
- (ii) $a \leq c$ and $d \leq b$.

Proof. Suppose that $f_0(x) \leq S(x + y) - P(y)$ for every $x \in M$ and $y \in E$. Then $f_0(w - z) \leq S(w - x_0) - P(z - x_0)$ for every $w, z \in M$, where $x = w - z$ and $y = z - x_0$. Consequently, from the linearity of f_0 , we obtain that $f_0(w) - S(w - x_0) \leq f_0(z) - P(z - x_0)$ for every $w, z \in M$. Hence, $\{f_0(w) - S(w - x_0) : w \in M\}$ is bounded above and has a least upper bound, a . Similarly, we can see that there is a $c = \inf_{z \in M} \{f_0(z) - P(z - x_0)\}$. Since $f_0(w - z) \leq S(w - x_0) - P(z - x_0)$ for every $w, z \in M$, it follows that $a \leq c$. If we let $x = w - z$ and $y = z + x_0$ in the inequality $f_0(x) \leq S(x + y) - P(y)$, we obtain that $-f_0(z) + P(z - x_0) \leq -f_0(w) + S(w + x_0)$ for every $w, z \in M$ and hence $d \leq b$. Conversely, assume that $a \leq c$ and $d \leq b$. Since $a \leq c$, it follows that $\sup_{w \in M} \{f_0(w) - S(w - x_0)\} \leq \inf_{z \in M} \{f_0(z) - P(z - x_0)\}$ and hence $f_0(w) - S(w - x_0) \leq f_0(z) - P(z - x_0)$ for every $w, z \in M$, and consequently $f_0(w - z) \leq S(w - x_0) - P(z - x_0)$. Thus, $f_0(x) \leq S(x + y) - P(y)$ for every $x = w - z \in M$ and $y = z - x_0 \in E$.

Remark 4.1. *Condition (i) in the Lemma 4.4 is equivalent to stating that $f_0(x) \leq T(x) = \inf_{y \in E} \{S(x + y) - P(y)\}$ for every $x \in M$. Also, condition (ii) in Lemma 4.4 is equivalent to the condition that $[a, b] \cap [d, c] \neq \emptyset$.*

Lemma 4.5. *Let $x_0 \notin M$ and let M_1 be the smallest linear subspace of E containing M and x_0 . Let f_0 be a linear functional on M such that $f_0(x) \leq S(x + y) - P(y)$ for every $x \in M$ and for every $y \in M_1$. Then there exists a linear functional L on M_1 such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in M_1$.*

Proof. From Lemma 4.1, we have $f_0(x) \leq S(x)$ for every $x \in M$. Therefore for arbitrary $y_1, y_2 \in M$ we have $f_0(y_1 + y_2) \leq S(y_1 + y_2)$. So, from the linearity of f_0 and the sublinearity of S we get $f_0(y_1) + f_0(y_2) = f_0(y_1 + y_2) \leq S(y_1 + y_2) \leq S(y_1 - x_0) + S(y_2 + x_0)$. Thus, we obtain:
 $f_0(y_1) - S(y_1 - x_0) \leq -f_0(y_2) + S(y_2 + x_0)$ (4.4).

By fixing y_2 and letting y_1 vary over M , we see that $\{f_0(y_1) - S(y_1 - x_0) : y_1 \in M\}$ is bounded above. Let $a = \sup_{y_1 \in M} \{f_0(y_1) - S(y_1 - x_0)\}$. By a similar argument, we can find $b = \inf_{y_2 \in M} \{-f_0(y_2) + S(y_2 + x_0)\}$. It is

clear from (4.4) that $a \leq b$. Similarly, from Lemma 4.1, we know that $P(x) \leq f_0(x)$ for every $x \in M$. Therefore, for arbitrary $y_1, y_2 \in M$ we have $P(y_1 + y_2) \leq f_0(y_1 + y_2)$. So, from the linearity of f_0 and from the super linearity of P , we get $P(y_1 + x_0) + P(y_2 - x_0) \leq P(y_1 + y_2) \leq f_0(y_1) + f_0(y_2)$. Thus, we have:

$$-f_0(y_1) + P(y_1 + x_0) \leq f_0(y_2) - P(y_2 - x_0) \quad (4.5).$$

By fixing y_2 and letting y_1 vary over M , we see that $\{-f_0(y_1) + P(y_1 + x_0) : y_1 \in M\}$ is bounded above. Let $d = \sup_{y_1 \in M} \{-f_0(y_1) + P(y_1 + x_0)\}$.

By a similar argument, there exists a $c = \inf_{y_2 \in M} \{f_0(y_2) - P(y_2 - x_0)\}$, and

from inequality (4.5) we see that $d \leq c$. By Remark 4.1, we note that $[a, b] \cap [d, c] \neq \emptyset$. Let $\xi \in [a, b] \cap [d, c]$. Note that, for every $x \in M_1$, there is a unique $y \in M$ and $\alpha \in \mathbb{R}$ such that $x = y + \alpha x_0$. We define a real valued functional L on M_1 by setting $L(x) = L(y + \alpha x_0) = f_0(y) + \alpha \xi$. We wish to verify that L is linear such that $L(x) = f_0(x)$ for every $x \in M$, and that $P(x) \leq L(x) \leq S(x)$ for every $x \in M_1$. First, we show that L is linear. Let $z_1, z_2 \in M_1$. Then there exist $y_1, y_2 \in M$ and $\alpha_1, \alpha_2 \in \mathbb{R}$ such that $z_1 = y_1 + \alpha_1 x_0$ and $z_2 = y_2 + \alpha_2 x_0$. Hence, for every $\alpha, \beta \in \mathbb{R}$ we get $L(\alpha z_1 + \beta z_2) = L((\alpha y_1 + \beta y_2) + (\alpha \alpha_1 + \beta \alpha_2) x_0) = f_0(\alpha y_1 + \beta y_2) + (\alpha \alpha_1 + \beta \alpha_2) \xi = \alpha(f_0(y_1) + \alpha_1 \xi) + \beta(f_0(y_2) + \alpha_2 \xi) = L(z_1) + L(z_2)$. Next, we wish to show that $L(x) = f_0(x)$ for every $x \in M$. If $x \in M$, then since $x = y + \alpha x_0$ for a unique choice of $y \in M$ and $\alpha \in \mathbb{R}$, it follows that that $x = y$ and $\alpha = 0$. Thus, $L(x) = f_0(x)$ for every $x \in M$, so $L|_M = f_0$. To show that $P(x) \leq L(x) \leq S(x)$ for every $x = y + \alpha x_0 \in M_1$, we consider three cases.

Case 1. $\alpha = 0$. We have already shown that $L(x) = f_0(x)$, and therefore $P(x) \leq L(x) \leq S(x)$, for every $x \in M$.

Case 2. $\alpha > 0$. Since $\xi \in [a, b] \cap [d, c]$, it follows that $d = \sup_{y \in M} \{-f_0(y) + P(y + x_0)\} \leq \xi \leq b = \inf_{y \in M} \{-f_0(y) + S(y + x_0)\}$. Thus, $-f_0(y) + P(y + x_0) \leq \xi \leq -f_0(y) + S(y + x_0)$ for every $y \in M$. Replacing y by $\frac{y}{\alpha}$, and multiplying the inequality by α , we get $P(y + \alpha x_0) \leq f_0(y) + \alpha \xi \leq$

$S(y + \alpha x_0)$ for every $y \in M$. Therefore $P(x) \leq L(x) \leq S(x)$ for every $x \in M_1$, where $\alpha > 0$.

Case 3. $\alpha < 0$. Let $\alpha = -t$, where $t > 0$. We know that

$$a = \sup_{y \in M} \{f_0(y) - S(y - x_0)\} \leq \xi \leq c = \inf_{y \in M} \{f_0(y) - P(y - x_0)\}$$

so $f_0(y) - S(y - x_0) \leq \xi \leq f_0(y) - P(y - x_0)$ for every $y \in M$. Thus, $-S(y - x_0) \leq -f_0(y) + \xi \leq -P(y - x_0)$ for every $y \in M$. Replacing y by $\frac{y}{t}$, and multiplying both sides by t , we get $-S(y - tx_0) \leq -f_0(y) + t\xi \leq -P(y - tx_0)$, so $-S(y + \alpha x_0) \leq -f_0(y) - \alpha\xi \leq -P(y + \alpha x_0)$. This means that $P(y + \alpha x_0) \leq f_0(y) + \alpha\xi \leq S(y + \alpha x_0)$ for every $y \in M$. Therefore, $P(x) \leq L(x) \leq S(x)$ for every $x \in M_1$. This completes the proof.

Theorem 4.2. *Let M and E_1 be two linear subspaces of E such that $M \subseteq E_1 \subseteq E$. Suppose that f_0 is a linear functional on M such that $f_0(x) \leq S(x+y) - P(y)$ for every $x \in M$ and $y \in E_1$. Then there exists a linear functional L on E_1 such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in E_1$.*

Proof. The proof follows directly from Lemma 4.5 and Zorn's Lemma (See [3], p. 134 for this argument).

It is worth pointing out that such an extension of a linear functional f_0 on M does not always exist on the whole space E . If (4.1) is not valid on the whole space E but is valid on some linear subspace E_1 containing M , then by using the above theorem we can extend f_0 to a linear functional L on E_1 so that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in E_1$, but it does not follow that we can extend f_0 to E .

Example 4.2. Let $S(x) = \sqrt{\sum_{i=1}^4 x_i^2}$ and $P(x) = -\sqrt{\sum_{i=1}^3 x_i^2} + x_4$ for every $x \in \mathbb{R}^4$. Let $M = \{(x_1, 0, 0, 0) : x_1 \in \mathbb{R}\}$ be a subspace of $\mathbb{R}^4$, and define a linear functional f_0 on M as $f_0(x) = x_1$ for every $x = (x_1, 0, 0, 0) \in M$. Then we show that there exists a linear functional L on a subspace $E_1 = \mathbb{R}^3$ such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in \mathbb{R}^3$, but there does not exist a linear functional L on $\mathbb{R}^4$ such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in \mathbb{R}^4$.

Solution. It is easy to verify that $S \in \text{Subl}(\mathbb{R}^4)$ and $P \in \text{Supl}(\mathbb{R}^4)$. First, we show that $f_0(x) \leq S(x+y) - P(y)$ for every $x \in M$ and for every $y \in M_1 = \{(x_1, x_2, x_3, 0) : x_1, x_2, x_3 \in \mathbb{R}\}$. Let $x = (x_1, 0, 0, 0) \in$

M and $y = (y_1, y_2, y_3, 0) \in M_1$. Then $f_0(x) = x_1$, $S(x + y) = \sqrt{(x_1 + y_1)^2 + y_2^2 + y_3^2}$ and $P(y) = -\sqrt{y_1^2 + y_2^2 + y_3^2}$. Therefore $f_0(x) = x_1 \leq |x_1| = |x_1| - |y_1| + |y_1| \leq |x_1 + y_1| + |y_1| \leq \sqrt{(x_1 + y_1)^2 + y_2^2 + y_3^2} + \sqrt{y_1^2 + y_2^2 + y_3^2} \leq S(x + y) - P(y)$. Hence, by Theorem 4.2, there exists a linear functional L_1 on M_1 such that $L_1|_M = f_0$ and $P(x) \leq L_1(x) \leq S(x)$ for every $x \in M_1$. Suppose there is a linear functional L on $\mathbb{R}^4$ such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in \mathbb{R}^4$. Then by Theorem 4.1 and Remark 4.1, it must follow that $f_0(x) \leq S(x + y) - P(y)$ for every $x \in M$ and for every $y \in \mathbb{R}^4$. But if we let $x = (10, 0, 0, 0) \in M$ and $y = (0, 0, 0, 1) \in \mathbb{R}^4$, then $f_0(x) = 10$, $S(x + y) = \sqrt{101}$, $P(y) = 1$ and consequently $S(x + y) - P(y) \cong 9.489 < 10 = f_0(x)$, which contradicts the assumption that $f_0(x) \leq S(x + y) - P(y)$.

Theorem 4.3. *Let f_0 be a linear functional on M such that $f_0(x) \leq S(x + y) - P(y)$ for every $x \in M$ and $x \in E$. Then there exists a linear functional L on E such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in E$.*

Proof. The theorem follows directly from Theorem 4.2, setting $E = E_1$.

We now move to our third proof of the Sandwich Theorem for Superlinear and Sublinear Functionals. Throughout the remainder of this paper, let $\Gamma_1 = \{L \in \text{Lin}(E) : L|_M = f_0 \text{ and } L(x) \leq S(x) \text{ for every } x \in E\}$ and let $\Gamma_2 = \{L \in \text{Lin}(E) : L|_M = f_0 \text{ and } P(x) \leq L(x) \text{ for every } x \in E\}$.

Remark 4.2. *Let $f_0 \in \text{Lin}(M)$, $P \in \text{Supl}(E)$, and $S \in \text{Subl}(E)$, so that $P(x) \leq f_0(x) \leq S(x)$ for every $x \in M$, and $P(x) \leq S(x)$ for every $x \in E$. Then by Theorem 3.1 and see Theorem 3.3, it follows that Γ_1 and Γ_2 are non-empty.*

Theorem 4.4. *A necessary and sufficient condition for $\Gamma_1 \cap \Gamma_2 \neq \emptyset$ is that $f_0(x) \leq T(x)$ for every $x \in M$, where $T(x) = \inf_{y \in E} \{S(x + y) - P(y)\}$.*

Proof. Suppose there is a linear functional $L \in \Gamma_1 \cap \Gamma_2$. Then $L|_M = f_0$, $L(x) \leq S(x)$ and $P(x) \leq L(x)$ for every $x \in E$. Since $P(u) \leq L(u)$, $P(-u) \leq L(-u) = -L(u)$, and $L(u) \leq -P(-u)$ for every $u \in E$. Therefore, from the linearity of L , we have $L(u + v) = L(u) + L(v) \leq S(v) - P(-u)$. This means that $L(x) \leq S(x - u) - P(-u)$, so $L(x) \leq$

$S(x + y) - P(y)$, where $x = u + v \in E$ and $y = -u \in E$. Since $L|_M = f_0$, it follows that $f_0(x) \leq S(x + y) - P(y)$ for every $x \in M$ and $y \in E$. Taking the infimum over all $y \in E$, $f_0(x) \leq T(x)$ for every $x \in M$. Conversely, assume that $f_0(x) \leq T(x)$ for every $x \in M$. It is easy to verify that $T \in \text{Subl}(E)$ (see Lemma 4.3). By lemma 4.1, $P(x) \leq S(x)$ for all $x \in E$ and $P(x) \leq f_0(x) \leq S(x)$ for all $x \in M$. By Theorem 3.3 there is an extension linear functional L on E such that $f_0(x) = L(x)$ for every $x \in M$ and $P(x) \leq L(x)$ for every $x \in E$. By lemma 4.3 $L(x) \leq T(x) \leq S(x)$ for every $x \in E$. Hence, it follows that $P(x) \leq L(x) \leq S(x)$ for every $x \in E$. Hence, $L \in \Gamma_2 \cap \Gamma_1$.

Theorem 4.5. *Let f_0 be a linear functional on M , where M is a subspace of E such that $P(x) \leq f_0(x) \leq S(x)$ for every $x \in M$, where $S \in \text{Subl}(E)$ and $P \in \text{Supl}(E)$. Then the necessary and sufficient condition that there exists a linear functional L on E such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in E$ is that $f_0(x) \leq T(x)$ for every $x \in M$.*

Proof. Since f_0 is a linear functional on M and $P(x) \leq f_0(x) \leq S(x)$ for every $x \in M$, then $P(x) \leq f_0(x)$ and $f_0(x) \leq S(x)$, then from Remark 4.2, we conclude that $\Gamma_1 \neq \emptyset$ and $\Gamma_2 \neq \emptyset$. Assume that L is a linear functional on E such that $L|_M = f_0$ and $P(x) \leq L(x) \leq S(x)$ for every $x \in E$. Then $\Gamma_1 \cap \Gamma_2 \neq \emptyset$. Hence from Theorem 4.4, we have $f_0(x) \leq T(x)$ for every $x \in M$. Conversely, let $f_0(x) \leq T(x)$ for every $x \in M$. Then $T \in \text{Subl}(E)$, and by Theorem 4.4 we obtain that $\Gamma_1 \cap \Gamma_2 \neq \emptyset$, and consequently there exists a linear functional L on E such that $L|_M = f_0$ and $P(x) \leq L(x)$ for every $x \in E$ and $L(x) \leq S(x)$ for every $x \in E$.

REFERENCES

1. A.T.Diab, N. Faried and Afaf Abou Elmatty, *Extending Linear Continuous Functionals with Preservation of Positivity*, Proc. Math. Phys. Soc. Egypt, 84, 2, p.p. 125-132, (2006).
2. J. L. Kelley, I. Namioka, and co-authors, *Linear Topological Spaces*, D.Van Nostrand Co., Inc., Princeton-Toronto-London-Melbourne(1963).
3. A. N. Kolmogorov and S. V. Fomin, *Introductory Real Analysis*, Translated and Edited by Richard A. Silverman, Dover Publications, Inc. New York (1970).
4. H. Konig, *Some Basic Theorems in Convex Analysis*, in "Optimization and operations Research", edited by B. Korte, North-Holland (1982).

5. E. Muhamadiev and A. T. Diab, *On extension of positive linear functionals*, Internat. J. Math., Math. Sci. 23, 1 (2000).
6. A. S. Okb-El-Bab, A. T. Diab and R. M. Boshnaa, *Generalization of the Hahn-Banach theorem*, Proceedings of International Conference on a theme " The actual problems of mathematics and computer science", Baku, 14-15.01.2009.
7. W. Rudin, *Functional Analysis*, Mc Graw-Hill, New York (1973).
8. S. Simons, *A new version of the Hahn-Banach theorem*, Arch. Math., 80, p.p. 630-646, (2003).
9. S. Simons, *From Hahn-Banach to Monotonicity*, Lecture Notes in Mathematics, 1693, Second Edition, Springer Science + Business Media B.V., 1998 Springer-Verlag, Berlin Heidelberg (2008).

DEPT. OF MATH., FACULTY OF SCIENCE, AIN SHAMS UNIVERSITY, CAIRO, EGYPT

DEPT. OF MATH., FACULTY OF SCIENCE, QATAR UNIVERSITY, DOHA, QATAR

DEPT. OF MATH., UTAH VALLEY UNIVERSITY, OREM, UTAH, USA

E-mail address: adel diab80@hotmail.com

E-mail address: snada@qu.edu.qa

E-mail address: fearnlda@uvu.edu